

THE STATA JOURNAL

Editor

H. Joseph Newton
Department of Statistics
Texas A & M University
College Station, Texas 77843
979-845-3142; FAX 979-845-3144
jnewton@stata-journal.com

Executive Editor

Nicholas J. Cox
Department of Geography
University of Durham
South Road
Durham City DH1 3LE UK
n.j.cox@stata-journal.com

Associate Editors

Christopher Baum
Boston College
Rino Bellocco
Karolinska Institutet
David Clayton
Cambridge Inst. for Medical Research
Mario A. Cleves
Univ. of Arkansas for Medical Sciences
William D. Dupont
Vanderbilt University
Charles Franklin
University of Wisconsin, Madison
Joanne M. Garrett
University of North Carolina
Allan Gregory
Queen's University
James Hardin
University of South Carolina
Stephen Jenkins
University of Essex
Ulrich Kohler
WZB, Berlin
Jens Lauritsen
Odense University Hospital

Stanley Lemeshow
Ohio State University
J. Scott Long
Indiana University
Thomas Lumley
University of Washington, Seattle
Roger Newson
King's College, London
Marcello Pagano
Harvard School of Public Health
Sophia Rabe-Hesketh
University of California, Berkeley
J. Patrick Royston
MRC Clinical Trials Unit, London
Philip Ryan
University of Adelaide
Mark E. Schaffer
Heriot-Watt University, Edinburgh
Jeroen Weesie
Utrecht University
Nicholas J. G. Winter
Cornell University
Jeffrey Wooldridge
Michigan State University

Stata Press Production Manager

Lisa Gilmore

Copyright Statement: The Stata Journal and the contents of the supporting files (programs, datasets, and help files) are copyright © by StataCorp LP. The contents of the supporting files (programs, datasets, and help files) may be copied or reproduced by any means whatsoever, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the Stata Journal.

The articles appearing in the Stata Journal may be copied or reproduced as printed copies, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the Stata Journal.

Written permission must be obtained from StataCorp if you wish to make electronic copies of the insertions. This precludes placing electronic copies of the Stata Journal, in whole or in part, on publicly accessible web sites, fileservers, or other locations where the copy may be accessed by anyone other than the subscriber.

Users of any of the software, ideas, data, or other materials published in the Stata Journal or the supporting files understand that such use is made without warranty of any kind, by either the Stata Journal, the author, or StataCorp. In particular, there is no warranty of fitness of purpose or merchantability, nor for special, incidental, or consequential damages such as loss of profits. The purpose of the Stata Journal is to promote free communication among Stata users.

The *Stata Journal* (ISSN 1536-867X) is a publication of Stata Press, and Stata is a registered trademark of StataCorp LP.

The Stata Journal publishes reviewed papers together with shorter notes or comments, regular columns, book reviews, and other material of interest to Stata users. Examples of the types of papers include 1) expository papers that link the use of Stata commands or programs to associated principles, such as those that will serve as tutorials for users first encountering a new field of statistics or a major new technique; 2) papers that go “beyond the Stata manual” in explaining key features or uses of Stata that are of interest to intermediate or advanced users of Stata; 3) papers that discuss new commands or Stata programs of interest either to a wide spectrum of users (e.g., in data management or graphics) or to some large segment of Stata users (e.g., in survey statistics, survival analysis, panel analysis, or limited dependent variable modeling); 4) papers analyzing the statistical properties of new or existing estimators and tests in Stata; 5) papers that could be of interest or usefulness to researchers, especially in fields that are of practical importance but are not often included in texts or other journals, such as the use of Stata in managing datasets, especially large datasets, with advice from hard-won experience; and 6) papers of interest to those teaching, including Stata with topics such as extended examples of techniques and interpretation of results, simulations of statistical concepts, and overviews of subject areas.

For more information on the Stata Journal, including information for authors, see the web page

<http://www.stata-journal.com>

Subscriptions are available from StataCorp, 4905 Lakeway Drive, College Station, Texas 77845, telephone 979-696-4600 or 800-STATA-PC, fax 979-696-4601, or online at

<http://www.stata.com/bookstore/sj.html>

Subscription rates:

Subscriptions mailed to US and Canadian addresses:

3-year subscription (includes printed and electronic copy)	\$153
2-year subscription (includes printed and electronic copy)	\$110
1-year subscription (includes printed and electronic copy)	\$ 59
1-year student subscription (includes printed and electronic copy)	\$ 35

Subscriptions mailed to other countries:

3-year subscription (includes printed and electronic copy)	\$225
2-year subscription (includes printed and electronic copy)	\$158
1-year subscription (includes printed and electronic copy)	\$ 83
1-year student subscription (includes printed and electronic copy)	\$ 59
3-year subscription (electronic only)	\$153

Back issues of the Stata Journal may be ordered online at

<http://www.stata.com/bookstore/sj.html>

The Stata Journal is published quarterly by the Stata Press, College Station, Texas, USA.

Address changes should be sent to the Stata Journal, StataCorp, 4905 Lakeway Drive, College Station TX 77845, USA, or email sj@stata.com.


Stata tip 15: Function graphs on the fly

Nicholas J. Cox
University of Durham, UK
n.j.cox@durham.ac.uk

[G] **graph twoway function** gives several examples of how function graphs may be drawn on the fly. The manual entry does not quite explain the full flexibility and versatility of the command. Here is a further advertisement on its behalf. To underline a key feature: you do not need to create variables following particular functions. The command handles all of that for you. We will look at two more examples.

A common simple need is to draw a circle. A trick with **twoway function** is to draw two half-circles, upper and lower, and combine them. If you are working in some scheme using color, you will usually also want to ensure that the two halves are shown in the same color. The **aspect()** option was explained in Cox (2004).

```
. twoway function sqrt(1 - x * x), ra(-1 1) ||  
> function -sqrt(1 - x * x), ra(-1 1) aspect(1)  
> legend(off) yla(, ang(h)) ytitle(, orient(horiz)) clp(solid)
```



MacKay (2003, 316) asserts that, if we transform beta distributions of variables P between 0 and 1 to the corresponding densities over $\text{logit } P = \ln[P/(1 - P)]$, then we find always pleasant bell-shaped densities. In contrast, densities over P may have singularities at $P = 0$ and $P = 1$. This is the kind of textbook statement that should provoke some play with friendly statistical graphics software.


To explore MacKay's assertion, we need a standard result on changing variables (see, for example, Evans and Rosenthal 2004, theorems 2.6.2 and 2.6.3). Suppose that P is an absolutely continuous random variable with density function f_P , h is a function that is differentiable and monotone, and $X = h(P)$. The density function of X is then

$$f_X(x) = \frac{f_P\{h^{-1}(x)\}}{|h'\{h^{-1}(x)\}|}$$

In our case, $h(P) = \text{logit } P$, so that $h^{-1}(X) = \exp(X)/\{1+\exp(X)\}$ and $h'\{h^{-1}(X)\} = \{1 + \exp(X)\}^2/\exp(X)$. In Stata terms, beta densities transformed to the logit scale are the product of `betaden(p)` or `betaden(invlogit(x))` and `exp(x)/(1+exp(x))^2`. The latter term may be recognized as a logistic density function, which always has a bell shape.

An example pair of original and transformed distributions is given by the commands below. To explore further in parameter space, you need only vary the parameters from 0.5 and 0.5 (and, if desired, to vary the range).

```
. twoway function betaden(0.5,0.5,x), ytitle(density) xtitle(p)
. twoway function betaden(0.5,0.5,invlogit(x)) * (exp(x) / (1 + exp(x))^2),
> ra(-10 10) ytitle(density) xtitle(logit p)
```



References

- Cox, N. J. 2004. Stata tip 12: Tuning the plot region aspect ratio. *Stata Journal* 4(3): 357–358.
- Evans, M. J. and J. S. Rosenthal. 2004. *Probability and Statistics: The Science of Uncertainty*. New York: W. H. Freeman.
- MacKay, D. J. C. 2003. *Information Theory, Inference, and Learning Algorithms*. Cambridge: Cambridge University Press. Also available at <http://www.inference.phy.cam.ac.uk/mackay/itprnn/book.html>.